

A Polynomial-Time Algorithm for the Tridiagonal and Hessenberg P-Matrix Linear Complementarity Problem

B. Gärtner and M. Sprecher*

December 2, 2011

Abstract

We give a polynomial-time dynamic programming algorithm for solving the linear complementarity problem with tridiagonal or, more generally, Hessenberg P-matrices. We briefly review three known tractable matrix classes and show that none of them contains all tridiagonal P-matrices.

1 Introduction

Given a matrix $M \in \mathbb{R}^{n \times n}$ and a vector $q \in \mathbb{R}^n$, the *linear complementarity problem* $\text{LCP}(M, q)$ is to find vectors $w, z \in \mathbb{R}^n$ such that

$$w - Mz = q, \quad w, z \geq 0, \quad w^T z = 0. \quad (1)$$

It is NP-complete in general to decide whether such vectors exist [2]. But if M is a *P-matrix* (meaning that all principal minors—determinants of principal submatrices—are positive), then there are unique solution vectors $\tilde{w}, \tilde{z}$ for every right-hand side q [10]. It is unknown whether these vectors can be found in polynomial time [7].

The matrix $M = (m_{ij})_{i,j=1}^n$ is *tridiagonal* if $m_{ij} = 0$ for $|j - i| > 1$. More generally, M is *lower Hessenberg* if $m_{ij} = 0$ for $j - i > 1$, and M is *upper Hessenberg* if M^T is lower Hessenberg; see Figure 1.

In this note we show that $\text{LCP}(M, q)$ can be solved in polynomial time if M is a lower (or upper) Hessenberg P-matrix. Polynomial-time results already




Figure 1: A tridiagonal matrix (left) and a lower Hessenberg matrix (right); the nonzero entries are enclosed in bold lines.

exist for other classes of matrices, most notably Z-matrices [1], hidden Z-matrices [6], and transposed hidden K-matrices [9]. Section 6 shows that none of these classes contains all tridiagonal P-matrices.

For the remainder of this note, we fix a P-matrix $M \in \mathbb{R}^{n \times n}$ and a vector $q \in \mathbb{R}^n$.

2 The optimal basis

For $B \subseteq [n] := \{1, 2, \dots, n\}$, we let $\overline{M}_B$ be the $n \times n$ matrix whose i th column is the i th column of $-M$ if $i \in B$, and the i th column of the $n \times n$ identity matrix I_n otherwise. $\overline{M}_B$ is invertible for every set B , a direct consequence of M having nonzero principal minors. We call B a *basis* and $\overline{M}_B$ the associated *basis matrix*.

The *complementary pair* $(w(B), z(B))$ associated with the basis B is defined by

$$w_i(B) := \begin{cases} 0 & \text{if } i \in B \\ (\overline{M}_B^{-1} q)_i & \text{if } i \notin B \end{cases} \quad (2)$$

*Institute of Theoretical Computer Science, ETH Zurich, CH-8092 Zurich, Switzerland (gaertner@inf.ethz.ch)

and

$$z_i(B) := \begin{cases} (\overline{M}_B^{-1} q)_i & \text{if } i \in B \\ 0 & \text{if } i \notin B \end{cases}, \quad (3)$$

for all $i \in [n]$.

Lemma 2.1. *For every basis $B \subseteq [n]$, the following two statements are equivalent.*

(i) *The pair $(w(B), z(B))$ solves $\text{LCP}(M, q)$, meaning that $w = w(B)$, $z = z(B)$ satisfy (1).*

(ii) $\overline{M}_B^{-1} q \geq 0$.

If both statements hold, B is called an *optimal basis* for $\text{LCP}(M, q)$.

Proof. As a consequence of (2) and (3), $w = w(B)$ and $z = z(B)$ already satisfy $w - Mz = q$ and $w^T z = 0$, for every B . Moreover, $w, z \geq 0$ if and only if $w(B), z(B) \geq 0$; this in turn is equivalent to $\overline{M}_B^{-1} q \geq 0$. $\square$

From now on, we assume w.l.o.g. that $\text{LCP}(M, q)$ is *nondegenerate*, meaning that $(\overline{M}_B^{-1} q)_i \neq 0$ for all $B \subseteq [n]$ and all $i \in [n]$. We can achieve this e.g. through a symbolic perturbation of q . In this case, we obtain the following

Lemma 2.2. *There is a unique optimal basis $\tilde{B}$ for $\text{LCP}(M, q)$.*

Proof. Let $\tilde{w}, \tilde{z}$ be solution vectors of $\text{LCP}(M, q)$, and set $\tilde{B} := \{i \in [n] : \tilde{w}_i = 0\}$. Since $\tilde{w}^T \tilde{z} = 0$, we have $\tilde{z}_i = 0$ if $i \in [n] \setminus \tilde{B}$. Hence, the vectors $\tilde{w}, \tilde{z}$ satisfy

$$q = (I_n \mid -M) \begin{pmatrix} \tilde{w} \\ \tilde{z} \end{pmatrix} = \overline{M}_{\tilde{B}} \begin{pmatrix} \tilde{w}_{[n] \setminus \tilde{B}} \\ \tilde{z}_{\tilde{B}} \end{pmatrix},$$

so $\tilde{w} = w(\tilde{B})$, $\tilde{z} = z(\tilde{B})$ follows. Hence, $\tilde{B}$ satisfies statement (i) in Lemma 2.1 and is therefore an optimal basis.

Uniqueness of $\tilde{w}, \tilde{z}$ [10] implies via Lemma 2.1 that $(w(B), z(B)) = (w(\tilde{B}), z(\tilde{B}))$ for every optimal basis B . But then (2) and (3) show that $(\overline{M}_B^{-1} q)_i = (\overline{M}_{\tilde{B}}^{-1} q)_i = 0$ for all $i \in \tilde{B} \oplus B$. Under nondegeneracy, there can be no such i , hence $B = \tilde{B}$. $\square$

3 Subproblems

For $K \subseteq [n]$, let M_{KK} be the principal submatrix of M consisting of all entries m_{ij} with $i, j \in K$. Furthermore, let q_K be the subvector of q consisting of all entries $q_i, i \in K$.

By definition, the submatrix M_{KK} is also a P-matrix, and $\text{LCP}(M_{KK}, q_K)$ is easily seen to inherit nondegeneracy from $\text{LCP}(M, q)$. Hence, Lemma 2.2 allows us to make the following

Definition 3.1. *For $k \in [n]$, $B(k) \subseteq [k]$ is the unique optimal basis of $\text{LCP}(M_{[k][k]}, q_{[k]})$.*

We also set $B(-1) = B(0) = \emptyset$.

4 The lower Hessenberg case

Let M be a lower Hessenberg matrix. Then we have the following

Theorem 4.1. *For every $k \in [n]$, there exists an index $\ell \in \{-1, 0, \dots, k-1\}$ such that*

$$B(k) = B(\ell) \cup \{\ell+2, \ell+3, \dots, k\}.$$

Proof. If $B(k) = [k]$, the statement holds with index $\ell = -1$. Otherwise, let $\ell \in \{0, 1, \dots, k-1\}$ be the largest index such that $\ell+1 \notin B(k)$. The matrix $M_{[k][k]}$ is lower Hessenberg as well, which implies that the basis matrix $\overline{M} := \overline{M}_{[k][k]B(k)}$ associated with $B(k)$ satisfies $\overline{m}_{ij} = 0$ if $i \leq \ell < j$; see Figure 2.




Figure 2: The basis matrix $\overline{M} = \overline{M}_{[k][k]B(k)}$ in the tridiagonal and lower Hessenberg case if $\ell+1 \notin B_k$. We have $\overline{M}_{[\ell][\ell]} = \overline{M}_{[\ell][\ell]B(k) \cap [\ell]}$.

As a consequence, the system of k equations

$$\overline{M}_{[k][k]B(k)} x = q_{[k]} \quad (4)$$

includes the ℓ equations

$$\overline{M_{[\ell][\ell]B(k) \cap [\ell]}} x_{[\ell]} = q_{[\ell]}. \quad (5)$$

Since $B(k)$ is the optimal basis of $\text{LCP}(M_{[k][k]}, q_{[k]})$, the unique solution $\tilde{x}$ of (4) satisfies $\tilde{x} \geq 0$; see Lemma 2.1. Vice versa, the unique partial solution $\tilde{x}_{[\ell]} \geq 0$ of subsystem (5) shows that $B(k) \cap [\ell] = B(\ell)$, the unique optimal basis of $\text{LCP}(M_{[\ell][\ell]}, q_{[\ell]})$. Together with the choice of ℓ , the statement of the theorem follows. $\square$

We remark that a variant of Theorem 4.1 for upper Hessenberg matrices can be obtained by considering *lower right* principal submatrices M_{KK} .

5 Polynomial-time algorithm

A *basis test* is a procedure to decide whether a given basis $B \subseteq [n]$ is optimal for $\text{LCP}(M, q)$. According to Lemma 2.1, a basis test can be implemented in polynomial time, using Gaussian elimination. In the sequel, we will therefore adopt the number of basis tests as a measure of algorithmic complexity. Here is our main result.

Theorem 5.1. *Let $M \in \mathbb{R}^{n \times n}$ be a lower Hessenberg P-matrix. The optimal basis $\tilde{B} = B(n)$ of $\text{LCP}(M, q)$ can be found with at most $\binom{n+1}{2}$ basis tests.*

Proof. We successively compute the optimal bases $B(-1), B(0), \dots, B(n)$, where $B(-1) = B(0) = \emptyset$. To determine $B(k), k > 0$, we simply test the $k+1$ candidates for $B(k)$ that are given by Theorem 4.1. In fact, we already know $B(k)$ after testing k of the candidates. This algorithm requires a total of $\sum_{k=1}^n k = \binom{n+1}{2}$ basis tests. $\square$

Using an $O(n^3)$ Gaussian elimination procedure, we obtain an $O(n^5)$ algorithm—this is certainly not best possible. Faster algorithms are available if M is a tridiagonal Z-matrix [5] or K-matrix [4, 3], but to our knowledge, the above algorithm is the first one to handle tridiagonal (and lower Hessenberg) P-matrices in polynomial time. The case of upper Hessenberg matrices is analogous, see the remark at the end of Section 4.

All upper and lower triangular P-matrices are hidden Z [12], meaning that linear complementarity problems with triangular P-matrices can be solved in polynomial time [6]. We can now also handle the “almost” triangular Hessenberg P-matrices. As we show next, there is a significant combinatorial difference between the two classes.

6 A tridiagonal example

Let us consider $\text{LCP}(M, q)$ with

$$M = \begin{pmatrix} 36 & -81 & 0 & 0 \\ 147 & 16 & -74 & 0 \\ 0 & 114 & 28 & 171 \\ 0 & 0 & -33 & 72 \end{pmatrix}, q = \begin{pmatrix} 1 \\ 1 \\ -1 \\ 1 \end{pmatrix}. \quad (6)$$

This linear complementarity problem was found by a computer search, with the goal of establishing Lemma 6.2 below. It can be checked that M is a tridiagonal P-matrix, but not a Z-matrix (a matrix with nonpositive off-diagonal entries). To show that some other known polynomial-time manageable matrix classes fail to contain all tridiagonal P-matrices, we need a new concept.

Definition 6.1. *Let $\mathcal{O}(M, q)$ be the digraph with vertex set $2^{[n]}$ and arc set*

$$\{(B, B \oplus \{i\}) : B \subseteq [n], i \in [n], (\overline{M_B}^{-1} q)_i < 0\}.$$

This digraph was first studied by Stickney & Watson [11]. Under nondegeneracy of $\text{LCP}(M, q)$, it has a unique sink that coincides with the optimal basis.

Lemma 6.2. *For M, q as in (6), $\mathcal{O}(M, q)$ contains the directed cycle $\{1, 2, 3, 4\} \rightarrow \{1, 2, 3\} \rightarrow \{1, 2\} \rightarrow \{2\} \rightarrow \emptyset \rightarrow \{3\} \rightarrow \{3, 4\} \rightarrow \{2, 3, 4\} \rightarrow \{1, 2, 3, 4\}$.*

We omit the elementary proof. This implies that M cannot be a hidden Z-matrix, since for such matrices, $\mathcal{O}(M, q)$ is the acyclic digraph of some geometric hypercube in $\mathbb{R}^n$, with edges directed by a linear function [6]. For the same reason, the tridiagonal P-matrix M^T cannot be the transpose of a hidden K-matrix (a hidden Z-matrix that is also a P-matrix) [9]. We remark that Morris has constructed a family of lower Hessenberg matrices $M \in \mathbb{R}^{n \times n}$ such that $\mathcal{O}(M, q)$ is highly cyclic for suitable $q \in \mathbb{R}^n$ [8].

7 Beyond Hessenberg matrices

It is natural to ask whether $\text{LCP}(M, q)$ can still be solved in polynomial time if M is a matrix of fixed bandwidth (number of nonzero diagonals), or fixed half-bandwidth (number of nonzero diagonals above or below the main diagonal); see Figure 3.




Figure 3: Matrices of bandwidth 5 (left) and right half-bandwidth 2 (right).

Let M be of fixed right half-bandwidth t . Generalizing Theorem 4.1, one can prove that there are only polynomially many candidates for $B(k)$, provided that $B(k)$ has a t -hole, meaning that $B(k)$ is disjoint from some contiguous t -element subset of $[k]$.

The only subset of $[k]$ without a 1-hole is the set $[k]$ itself, and this is why the lower Hessenberg case $t = 1$ is easy. But there is already an exponential number of subsets of $[k]$ without a 2-hole. Hence, the above approach fails for $t \geq 2$. It remains open whether there is another polynomial-time algorithm in the case of fixed (right) bandwidth.

References

- [1] R. Chandrasekaran. A special case of the complementary pivot problem. *Opsearch*, 7:263–268, 1970.
- [2] S.-J. Chung. NP-completeness of the linear complementarity problem. *J. Optim. Theory Appl.*, 60(3):393–399, 1989.
- [3] R. W. Cottle and R. S. Sacher. On the solution of large, structured linear complementarity problems: The tridiagonal case. *Applied Mathematics & Optimization*, 3:321–340, 1976.
- [4] C. W. Cryer. The efficient solution of linear complementarity problems for tridiagonal Minkowski matrices. *ACM Trans. Math. Softw.*, 9:199–214, June 1983.
- [5] S. Gupta. Linear time algorithm for linear complementarity problems with tridiagonal Z matrices. Technical report, Pennsylvania State University, Dept. of Computer Science, 1987.
- [6] O. L. Mangasarian. Linear complementarity problems solvable by a single linear program. *Mathematical Programming*, 10:263–270, 1976.
- [7] N. Megiddo. A note on the complexity of P-matrix LCP and computing an equilibrium. RJ 6439, IBM Research, Almaden Research Center, 650 Harry Road, San Jose, California, 1988.
- [8] W. D. Morris jr. Randomized pivot algorithms for P-matrix linear complementarity problems. *Mathematical Programming*, 92:285–296, 2002.
- [9] J. S. Pang and R. Chandrasekaran. Linear complementarity problems solvable by a polynomially bounded pivoting algorithm. In Richard W. Cottle et al., editors, *Mathematical Programming Essays in Honor of George B. Dantzig Part II*, volume 25 of *Mathematical Programming Studies*, pages 13–27. Springer Berlin Heidelberg, 1985.
- [10] H. Samelson, R. M. Thrall, and O. Wesler. A partition theorem for Euclidean n -space. *Proc. Amer. Math. Soc.*, 9:805–807, 1958.
- [11] A. Stickney and L. Watson. Digraph models of Bard-type algorithms for the linear complementarity problem. *Math. Oper. Res.*, 3(4):322–333, 1978.
- [12] M. J. Tsatsomeros. Generating and detecting matrices with positive principal minors. *Asian Information-Science-Life*, 1(2):115–132, 2002.